

AD-A120 020

AIR FORCE AEROSPACE MEDICAL RESEARCH LAB WRIGHT-PATT--ETC F/G 6/16  
CARTILAGE OF THE INTERVERTEBRAL DISC ENG-PLATE, A HISTOLOGICAL,--ETC(U)  
AUG 82 N S NUSSBAUM  
AFAMRL-TR-81-122

UNCLASSIFIED

NL

1 of 1

AD A  
120020

END  
DATE  
FILMED  
11-88  
DTIC

AD A120020

AFAMRL-TR-81-122



**CARTILAGE OF THE INTERVERTEBRAL DISC END-PLATE  
A HISTOLOGICAL, HISTOCHEMICAL, AND FINE STRUCTURE STUDY**

*NOEL S. NUSSBAUM*

AUGUST 1981



DTIC FILE COPY

Approved for public release; distribution unlimited

AEROSPACE MEDICAL RESEARCH LABORATORY  
AEROSPACE MEDICAL DIVISION  
AIR FORCE SYSTEMS COMMAND  
WRIGHT-PATTERSON AIR FORCE BASE, OHIO 45433

92 10 06 017

## NOTICES

When US Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from Air Force Aerospace Medical Research Laboratory. Additional copies may be purchased from:

National Technical Information Service  
5285 Port Royal Road  
Springfield, Virginia 22161

Federal Government agencies and their contractors registered with Defense Technical Information Center should direct requests for copies of this report to:

Defense Technical Information Center  
Cameron Station  
Alexandria, Virginia 22314

## TECHNICAL REVIEW AND APPROVAL


AFAMRL-TR-81-122

The experiments reported herein were conducted according to the "Guide for the Care and Use of Laboratory Animals," Institute of Laboratory Animal Resources, National Research Council.

This report has been reviewed by the Office of Public Affairs (PA) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

**FOR THE COMMANDER**



HENNING E. VON GIERKE, Dr Ing  
Director  
Biodynamics and Bioengineering Division  
Air Force Aerospace Medical Research Laboratory

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFAMRL-TR-81-122	2. GOVT ACCESSION NO. AD-A320200	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CARTILAGE OF THE INTERVERTEBRAL DISC END-PLATE A histological, histochemical, and fine structure study	5. TYPE OF REPORT & PERIOD COVERED Technical Report 1 July 1980-30 September 81	
7. AUTHOR(s) Noel S. Nussbaum	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Air Force Aerospace Medical Research Laboratory Aerospace Medical Division, Air Force Systems Command, Wright-Patterson AFB, Ohio 45433	8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 7231-14-16 62202F	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE August 1982	
	13. NUMBER OF PAGES 15	
	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Vertebral Disc Degeneration      Cartilage End Plate Cartilage Ultrastructure Scanning Electron Microscopy Proteoglycan Histochemistry		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The cartilaginous end plates define the anatomical boundaries of the disc, serve as an area of contact with the vertebral body, and act as a semi-permeable membrane. The anatomic relationships that exist between the end plate and immediately adjacent structures can be better understood if ultrastructure is interpreted in conjunction with established histology. Undecalcified sections of Rhesus ( <i>Macaca mulatta</i> ) spinal segments were prepared, for both light and scanning electron microscopy, from 17 adolescent and young		

adults with radiographically normal spines. Alternate thick (100  $\mu$ m) and thin (12  $\mu$ m) sections were cut at 0.5 mm intervals through quick-frozen tissues, embedded in sodium carboxymethyl cellulose, on a LKB 2250 PMV Cryomicrotome. Thin sections were stained for histological and histochemical examination and thick sections were examined with an Etec Auto Scan electron microscope. End plate cartilage from mature spines demonstrate a micro-heterogeneous structure. Strut-like inter-territorial regions (ca. 50  $\mu$ m wide),--impregnated with calcium salts (von Kossa),--separate and define inter-connecting cellular and pericellular domains. The proteoglycan content of the pericellular matrix stains more intensely (Safranin O) than does the inter-territorial region. Structural and histochemical data suggest the presence of a cell-mediated diffusion pathway allowing for trans-endplate movement of disc nutrients. This interpretation contrasts with the classical view of the end plate as a passive diffusion pathway but may explain the observation that the bone-disc interface is only partially permeable in contrast to the freely permeable peripheral annulus. Degenerative alterations in this system could be expected to compromise disc function. A better understanding of these relationships is vital for the development of treatment strategies designed to alleviate and/or correct the effects of acute and chronic disc degeneration.

Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



## **PREFACE**

This work was performed while the author was a Visiting Professor at the Biodynamic Effects Branch (BBD), Biodynamics and Bioengineering Division, Air Force Aerospace Medical Research Laboratory. The assistance of K. C. Smith (BBD), R. Nieser and D. Mattie (Toxic Hazards Division, Pathology Branch) is appreciated.

## TABLE OF CONTENTS

	Page
INTRODUCTION .....	3
OBJECTIVE .....	3
MATERIALS AND METHODS .....	4
RESULTS .....	5
DISCUSSION .....	12
REFERENCE .....	13

## LIST OF ILLUSTRATIONS

### Figure

1 View of microtome within LKB Cryochest. Specimen block is to left of knifeholder. Frontal sections are being cut. 1/6X (Total magnification given for all figures)	6
2 Example of a reference photograph taken every 0.5 mm as specimen is sectioned. In this case the animal is identified as rhesus 495 and the section is 4.5 mm into the vertebral centrum in the sagital plane. 3/4X	6
3 Sagital section of rhesus T <sup>2</sup> -T <sup>3</sup> Intervertebral Disc. Trabecular bone of vertebral body encloses marrow spaces. Cartilagenous end-plates are separated by growth cartilage from the adjacent spongy bone. Mallory, 7.5X	7
4 Frontal Section of rhesus T <sup>6</sup> -T <sup>7</sup> Disc. Vascularization of inferior end-plate is continuous but is restricted to lateral ends of superior. Mallory, 7.5X	7
5 Low power photomicrograph of anterior margin of inferior end-plate. Vascular buds are surrounded by ossifying matrix. Collagen bundles of annulus fibrosus above terminate in cartilage of end-plate. Light staining growth cartilage is located between end-plate and subchondral bone of vertebral centrum below. 50X	8
6 Higher power view of disc-bone junction from young animal. Note absence of vascular buds and ossification centers in end-plate cartilage. Other features the same as fig. 5. 100X	8
7 Disc-bone junction below nucleus pulposus. Termination of annular collagen bundles associated with narrow zone of end-plate cartilage. Cartilage exhibits mosaic of staining intensities with darker territorial regions surrounding cells and interterritorial zone appearing lighter. Mallory stain. 150X	9
8 Higher power view of end-plate cartilage demonstrating mosaic of aniline blue staining associated with territorial (T) and interterritorial (I) regions. Mallory stain 450X	9
9 Positive histochemical reaction for proteoglycans in territorial zone. Alcian blue stain. 900X	10
10 Left. Positive von Kossa reaction demonstrating calcium salts in interterritorial zone. Right. Positive alcian blue reaction for territorial regions. 200X	10
11 Scanning electron micrograph (SEM) of end-plate cartilage. Cellular territorial (T) domains appear depressed between interterritorial (I) calcified regions. Compare with fig. 9. 410X	11
12 SEM of cellular elements, territorial zone and adjacent calcified interterritorial matrix. 1400X	11

## INTRODUCTION

Age-dependent fibrotic degeneration of the intervertebral disc seriously compromises energy dissipation within the compression loaded spine. The loss of viscoelasticity associated with fibrosis of the nucleus pulposus has been identified with impaired nutritive flow from adjacent subchondral bone. Development of a dynamic model of the human spine is in turn dependent upon quantification of relevant parameters. The influence of age-driven variables affecting these parameters must also be estimated.

## OBJECTIVE

The objectives of this study are twofold. The first is to describe the histological, histochemical, and ultrastructural features of the cartilagenous end plates of the intervertebral discs of the rhesus monkey (*Macaca mulatta*) as derived from examination of undecalcified sections. The second is to relate this structural information to current theories of disc nutrition, and age-dependent fibrotic degeneration.



## MATERIALS AND METHOD

Vertebral segments comprising the T<sup>7</sup> - T<sup>11</sup> region were isolated from fresh quick-frozen spines of female rhesus monkeys ranging in weight from 12.5 to 21.5 pounds. Five spines from fresh material and an additional 12 long-term frozen spines were examined. Frontal, sagittal and horizontal sections were cut with the LKB 2250 PMV cryomicrotome. Frozen, undecalcified specimens were embedded in carboxymethyl cellulose within a frame, and frozen in hexane cooled with dry ice (-75°C). The frozen block was then sectioned at 0.5 mm intervals (Fig. 1). Photographs of the block face (Fig. 2), a series of thin sections at 10-12 micrometers, and a thick section at 100 micrometers were collected at each 0.5 mm interval.

Sections were collected on transparent tape. Thin sections were fixed immediately in:

95% Ethyl alcohol	70 ml
10% Formalin	29 ml
Acetic acid (glacial)	1 ml

Fixation for 2 minutes was followed by a distilled water rinse and then the sections were allowed to air dry completely before staining. Adjacent sections were stained with H-E, Mallory's connective tissue stain, von Kossa for calcium salts, and Alcian blue-light green for proteoglycans (Humason, 1972). Microphotographs were prepared with a Zeiss Photomicroscope.

Thick sections for scanning electron microscopy were dehydrated overnight within the cryochamber. The sections were then removed to room temperature and allowed to dry completely (30 min. - 1 hr.). Appropriate regions were trimmed from the tape-mounted tissues and attached, unfixed, to SEM specimen stubs. Gold-palladium sputter coated specimens were examined and photographed with an Eteck Autoscan electron microscope.

## RESULTS

The processes of growth and maturation, represented in the disc-bone interface, create a more rigid union between these elements. The disc of the young individual is separated from the underlying vertebral cancellous bone by a relatively thick and discrete cartilage layer. Secondary centers of ossification associated with vascular invasion of this region, eventually form a continuous bony plate immediately below the annulus fibrosus, obliterating all but a thin remnant of the original cartilage structure. This process begins at the anterior-lateral margin of the inferior end-plate of each disc (Fig. 3). Fusion of discrete centers across the mid-line completes the process (Fig. 4).

While bony replacement of the cartilage plate has been initiated at the anterior margin, growth cartilage of the underlying cancellous bone is still active (Fig. 5) and many areas of contact between the end-plate and the underlying marrow spaces are retained (Fig. 6).

A feature, previously unreported, of this end-plate cartilage, is its mosaic appearance (Fig. 7). This is a reflection of denser territorial staining in comparison with the lighter staining interterritorial regions. This feature is present regardless of staining technique (Fig. 8).

Histochemical evaluation with Alcian blue demonstrates that the territorial regions are enriched with proteoglycans (Fig. 9). The interterritorial zone is conversely deficient in proteoglycan but does contain amorphous calcium salts (Fig. 10).

A similar picture emerges when this area is examined by scanning electron microscopy. A clear separation of territorial and interterritorial regions is visualized (Fig. 11). The textural elements of the fibrous and nonfibrous matrix elements are easily distinguished within these zones (Fig. 12). Cellular components are surrounded by a looser matrix than that of the more tightly packed interterritorial zone.




Fig. 1 View of microtome within LKB Cryochest. Specimen block is to left of knifeholder. Frontal sections are being cut. 1 6X (Total magnification given for all figures)




Fig. 2 Example of a reference photograph taken every 0.5 mm as specimen is sectioned. In this case the animal is identified as rhesus 495 and the section is 4.5 mm into the vertebral centrum in the sagittal plane. 3/4X




Fig. 3 Sagital section of rhesus T<sub>2</sub>-T<sub>3</sub> Intervertebral Disc. Trabecular bone of vertebral body encloses marrow spaces. Cartilagenous end-plates are separated by growth cartilage from the adjacent spongy bone. Mallory, 7.5X



Fig. 4 Frontal Section of rhesus T<sub>6</sub>-T<sub>7</sub> Disc. Vascularization of inferior end-plate is continuous but is restricted to lateral ends of superior. Mallory, 7.5X




Fig. 5 Low-power photomicrograph of anterior margin of inferior end-plate. Vascular buds are surrounded by ossifying matrix. Collagen bundles of annulus fibrosus above terminate in cartilage of end-plate. Light staining growth cartilage is located between end-plate and subchondral bone of vertebral centrum below. 50X




Fig. 6 Higher power view of disc-bone junction from young animal. Note absence of vascular buds and ossification in end-plate cartilage. Other features the same as fig. 5. 100X




Fig. 7 Disc-bone junction below nucleus pulposus. Termination of annular collagen bundles associated with narrow zone of end-plate cartilage. Cartilage exhibits mosaic of staining intensities with darker territorial regions surrounding cells and interterritorial zone appearing lighter. Mallory stain. 150X



Fig. 8 Higher power view of end-plate cartilage demonstrating mosaic of aniline blue staining associated with territorial (T) and interterritorial (I) regions. Mallory stain 450X




Fig. 9 Positive histochemical reaction for proteoglycans in territorial zone. Alcian blue stain. 900X



Fig. 10 Left. Positive von Kossa reaction demonstrating calcium salts in interterritorial zone. Right. Positive alcian blue reaction for territorial regions. 200X




Fig. 11 Scanning electron micrograph (SEM) of end-plate cartilage. Cellular territorial (T) domains appear depressed between interterritorial (I) calcified regions. Compare with fig. 9. 410X




Fig. 12 SEM of cellular elements, territorial zone and adjacent calcified interterritorial matrix. 1400X



## DISCUSSION

The intervertebral disc is specially adapted to resist compression loading while allowing spinal movement. It has been proposed that clinical disorders of the spine are the result of disc degeneration (Eyring, 1969). The mechanical properties of the disc are determined by the biochemical and biophysical state of the matrix components comprising the major substance of the nucleus pulposus and the noncollagenous elements of the annulus fibrosus (Muir, 1979; Lyons et al., 1981). This proteoglycan-enriched matrix is characterized by cell-mediated turnover, though the observed cell population is sparse (Nachemson et al., 1970).

After the age of 30 in humans, gradual changes in the biochemical and structural character of the discs become evident (Coventry, 1969). It has been suggested that these degenerative changes represent a form of accelerated aging (Harris & Macnab, 1954) and are responsible in part for the compromised biomechanics of the older spine. The nutritive needs of the cell populations within the nucleus and annulus compartments have been estimated (Maroudas et al., 1968) and it has been proposed that closure of the subchondral diffusion pathway, a consequence of ossification of the cartilaginous end-plate, is a determining factor in the ensuing cell death and matrix degeneration (Urban et al., 1977).

Histological examination of an aging series of human discs supported the significance of reduced trans end-plate diffusion in the etiology of disc degeneration (Nachemson et al., 1970). These and related studies consider the end-plates to be composed of hyaline cartilage and thus homologues of articular cartilage. Accordingly, diffusion and compression induced bulk flow are modeled upon data derived from examination of synovial joints (Brodin, 1955).

The results of this study in rhesus indicate, that while present, the cartilage of the end-plate is quite different in structure and presumably biomechanical function from that of articular cartilage. The identification of a heterogeneous extra-cellular matrix composed of calcified struts separating proteoglycan enriched territorial zones, suggests a unique relationship allowing for maximal linear strength coincident with prolonged cell viability. As ossification of this end-plate progresses, vascular invasion of the calcified matrix reestablishes a capillary bed immediately adjacent to the disc elements. Whether this vascular association is adequate to maintain cell function within the disc is dependent on structural and functional factors that are yet to be determined. Examination of undecalcified human disc is necessary in order to relate these observations to the clinical arena.

## References

- Brodin, H. 1955. "Paths of nutrition in articular cartilage and intervertebral discs." Acta Orthop. Scandinav. 24:177-183.
- Coventry, M. B. 1969. "Anatomy of the intervertebral disc." Clin. Orthop. 67:9-15.
- Eyring, E. J. 1969. "The biochemistry and physiology of the intervertebral disk." Clin. Orthop. 67:16-28.
- Harris, R. I. & Macnab, I. 1954. "Structural changes in the lumbar intervertebral discs." J. Bone & Joint Surg. 36B:304-322.
- Humason, G. L. 1972. Animal Tissue Techniques, 3rd ed. W. H. Freeman & Co., San Francisco.
- Lyons, G., Eisenstein, S. M. & Sweet, M. B. E. 1981. "Biochemical changes in intervertebral disc degeneration." Biochem. Biophys. Acta 673:443-453.
- Maroudas, A. Bullough, P., Swanson, S.A. & Freeman, M.A.R. 1968. "The permeability of articular cartilage." J. Bone & Joint Surg. 50B:166-177.
- Muir, I. H. M. 1979. Biochemistry, in: Adult Articular Cartilage, 2nd ed., Freeman, M.A.R. (ed.), Pitman Medical Publishing Co., Ltd., Turnbridge Wells, England.
- Nachemson, A., Lewis, T., Maroudas, A. & Freeman, M.A.R. 1970. "In Vitro diffusion of dye through the end-plates and the annulus fibrosus of human lumbar intervertebral discs." Acta Orthop. Scandinav. 41:589-607.
- Urban, J.P.G., Holm, S., Maroudas, A. & Nachemson, A. 1977. "Nutrition of the intervertebral disk." Clin. Orthop. 129:101-114.

END

DATE  
FILMED

11-82

DTIC